



Computer Aided Designing of New Non Steroidal Anti Inflammatory Drugs (NSAIDs) considering Ibuprofen as a Lead Compound followed by suggestion of a Good Synthetic Route.

Vijay Masand^a, A.A.Shaikh^b, Y.I Shaikh^b, G.M.Nazeruddin^{b*}

a) Dept. of Chemistry, Vidyabharti College, Camp, Amravati, Maharashtra, India

b) Dept. of Chemistry (PG and Research Centre) Poona College of Arts, Science and Commerce, Camp, Pune-411001, Maharashtra, India.

e-mail: gmnazeruddin@yahoo.co.in

Abstract:

Discovery of a new drug is a very difficult task. Pharmaceutical and biotechnology companies need to make huge investments in the discovery of a single drug. Most pharmaceutical or biotechnology companies claim that it costs anywhere between \$800 million to \$900 million and a time span of twelve to fifteen years to discover a new drug. In silico-chemico-biological approach computer plays very important role in discovery of new drug, not only it can save money but also time, and are believed to offer means of improved efficiency for the industry. Quantitative structure–activity relationship/Quantitative structure property relationship (QSAR/QSPR) methods represent an attempt to correlate structural and/or physical properties and descriptors of compounds with biological activities. we have designed various new non steroidal anti-inflammatory drugs (NSAIDs) and calculated various physical properties and molecular descriptors like log P, Dipole moment, Heat of formation, Ionization Potential, HOMO (Highest occupied molecular orbital) and LUMO (lowest unoccupied molecular orbital) energies and pk_a by using different software such as VegaZZ, Mopac, ADME and ChemDraw etc. and compared them with the lead compound, Ibuprofen. The newly designed compounds, which are having the comparable properties with the lead compound, are selected and a good synthetic route is suggested by following rules of green chemistry.

Key words: Designing, α -aryl Propanoic acid derivatives (NSAID), Physicochemical properties.




Introduction:

In the early 1960s, Corwin Hansch[1] extended the concept of Linear –free energy relations (LFER) to describe the effectiveness of biologically-active molecule. This represented efforts to quantitatively relate the structure of a compound to its activity and resulting equations were aptly named quantitative structure activity relationships (QSAR). Today, these equations are also called quantitatively structure property relationship (QSPR). Generating useful Hansch equation can be very challenging and even a good Hansch equation will not give perfect predication of activity. For this reason new methods have somewhat replaced the traditional Hansch analysis. In the late 1980s and early 1990s combinatorial chemistry emergent diminished the importance of QSAR. Since large libraries of compounds bearing varying substituents could be easily prepared, being able to predict activity was no longer necessary, simply make all the compounds one can imagine and test them in high-throughput screens.

Since the middle 1990s, a technique called comparative molecular field Analysis (COMFA)[2] has emerged. This method uses highly complicated statistical analysis with large number of variable to correlate practical molecular properties to activity. However, Drug designing is still a big challenge. Pharmaceutical and biotechnology companies need to make huge investments in the discovery of a single drug. Most pharmaceutical or biotechnology companies claim that it costs anywhere between \$800 million to \$900 million and a time span of twelve to fifteen years to discover a new drug. In silico-chemico-biological approach[3] computer plays very important role in discovery of new dug, not only it can save money but also time, and are believed to offer means of improved efficiency for the industry. Quantitative structure–activity relationship/Quantitative structure property relationship (QSAR/QSPR) methods represent an attempt to correlate structural and/or physical properties and descriptors of compounds with biological activities. Pharmaceutical and biotechnology companies need to make huge investments in the discovery of a single drug. Most pharmaceutical or biotechnology companies claim that it costs anywhere between \$800 million to \$900 million and a time span of twelve to fifteen years to discover a new drug. In silico-chemico-biological approach computer plays very important role in discovery of new dug, not only it can save money but also time, and are

believed to offer means of improved efficiency for the industry. Quantitative structure–activity relationship/Quantitative structure property relationship (QSAR/QSPR) methods represent an attempt to correlate structural and/or physical properties and descriptors of compounds with biological activities. This is continuation of our earlier work[4] We wish to report here a simple hypothesis, in which certain physical properties such as Heat of formation, Dipole moment, Ionization potential, HOMO, LUMO energies, should be calculated for newly design α -aryl propanoic acid derivatives and compared with the lead compound Ibuprofen. The derivatives which are having comparable properties are selected and a good synthetic route is suggested by following rules of green chemistry. They may be synthesized by a multicomponent reaction (Beignelli reaction)[5], a condensation of thiourea 5-(4-R-phenyl)-5-oxopentanoic acid and substituted benzaldehyde (scheme-1)



2a, R=CH₃; **3a**, R= H; **4a**, R=Cl; **5a**, R=OCH₃ [R₁=H]

2b, R₁=2-OH, 2-Cl, 2-NO₂, 2-OCH₃ [R=CH₃]

3b, R₁=2-OH, 2-Cl, 2-NO₂, 2-OCH₃ [R=H]

4b, R₁=2-OH, 2-Cl, 2-NO₂, 2-OCH₃ [R=Cl]

5b, R₁=2-OH, 2-Cl, 2-NO₂, 2-OCH₃ [R=OCH₃]

Computational Work and Calculation:

There are number of software such as Winmopac /Hyper chem. /Dragon etc. are available in the market through them not only various new derivatives of α -aryl propanoic acid can be designed but also various molecular descriptors[6] can be calculated. Similarly the same calculation can be done for the lead compound Ibuprofen followed by comparison of these values. Various



compounds were designed and physical properties were calculated, However for comparison twenty compounds, calculations are depicted in Table-1.

Table-1:

Sr. No.	Mol Id	MLOGP	ALOGP	AM1 HOMO	AM1 LUMO	AM1 IP	AM1 dipole
1.	2a	2.796	3.026	-8.907	-0.207	8.907	2.652
2.	2b-2-OH	2.292	2.758	-8.79	-0.111	8.79	3.048
3.	2b-2-Cl	3.287	3.69	-8.934	-0.235	8.934	2.418
4.	2b-2-NO₂	2.882	2.92	-9.006	-0.782	9.006	5.491
5.	2b-2-OCH₃	2.515	3.01	-8.7	-0.085	8.7	2.338
6.	3a	2.568	2.54	-8.916	-0.237	8.916	2.713
7.	3b-2-OH	2.064	2.272	-8.872	-0.201	8.872	3.995
8.	3b-2-Cl	3.064	3.204	-9.008	-0.183	9.008	2.43
9.	3b-2-NO₂	2.655	2.434	-9.038	-0.739	9.038	5.035
10.	3b-2-OCH₃	2.292	2.523	-8.731	-0.079	8.731	2.161
11.	4a	3.064	3.204	-9.032	-0.513	9.032	2.116
12.	4b-2-OH	2.56	2.937	-8.971	-0.492	8.971	3.729
13.	4b-2-Cl	3.555	3.869	-9.055	-0.534	9.055	1.808
14.	4b-2-NO₂	3.15	3.098	-9.133	-0.801	9.133	5.157
15.	4b-2-OCH₃	2.783	3.188	-8.857	-0.375	8.857	1.451
16.	5a	2.292	2.523	-8.885	-0.162	8.885	3.643
17.	5b-2-OH	1.8	2.256	-8.806	-0.151	8.806	5.249



18.	5b-2-Cl	2.783	3.188	-8.911	-0.197	8.911	2.717
19.	5b-2-NO₂	2.411	2.418	-8.994	-0.862	8.994	4.688
20.	5b-2-OCH₃	2.023	2.507	-8.75	-0.101	8.75	2.933
21.	Ibuprofen	3.23 100	3.582 100	-9.398 100	0.182 100	9.398 100	2.229 100

Results and discussions:

After comparing the various values of MLOGP, ALOGP, Austin Model 1 HOMO, Austin Model 1 LOMO, Austin Model 1 IP, Austin Model 1 dipole, the methoxy methyl and chloro derivatives are having quite good comparable properties with Ibuprofen. So they can be synthesized by Biginelli reaction. A condensation of thiourea 5-(4-R-phenyl)-5-oxopentanoic acid and substituted benzaldehyde.

Conclusions:

This is very simple hypothesis based on molecular descriptor and physico-chemical parameters to predict the biologically active compounds, they may have comparable activity to lead compound.

Acknowledgement

Authors are thankful to Anjuman Khairul Islam Trust, Mumbai for financial assistance

References:

1. Hansch, C, Muir, R M, Fujita, T, Maloney, P P, Geiger, F, and Streich, M J Am Chem Soc.; **1963**, 85, 2817. The Correlation of Biological Activity of Plant Growth Regulators and Chloromycetin Derivatives with Hammett Constants and Partition Coefficients
2. Cramer R.D.; Patterson D.E and Bunce J.D.; J. Am. Chem. Soc. **1988**; 110; 5959. Comparative molecular field analysis (CoMFA). 1. Effect of shape on binding of steroids to carrier proteins



3. Pozzan A.; *Curr PharmDes.* **2006**, *12*, 2099.
4. Hassan A. Osman, Sana Kauser, M.S. Pandharpatte; G.M. Nazeruddin *Archives Des Sciences*, **2013**, *66*, 503.
5. P. Biginelli, *Gazz. chim. Ital.*, **1893**, *23*, 360.
6. Todeschini R. and Consonni V.; "Handbook of Molecular Descriptor" Wiley-vch, **2000**