



COMPARATIVE EVALUATION FOR IN VITRO ANTIOXIDANT ACTIVITY OF *GYMNOSPORA MONTANA* LEAVES AND STEM

B. N. Gawade

Dept. of Chemistry, A. D. College, Kada. Tq. Ashti. Dist. Beed-414 202 (M.S.) India.

ABSTRACT

Gymnospora Montana occurring throughout the arid, dry areas of India and is traditionally claimed to be useful in various ailments. A detailed review of literature afforded no more information on the in vitro antioxidant activity of the same plant. The main objective of this study was to evaluate the free radical scavenging activity of *G.Montanaleaves* and stem by using DPPH radical scavenging method. For the present study ascorbic acid was used as reference standard and it shows IC₅₀ value 25 $\mu\text{g/ml}$. The extracts were prepared only in ethanol. The ethanol extracts of leaves and stem exhibited DPPH scavenging activity with IC₅₀ value 34 $\mu\text{g/ml}$ and 36 $\mu\text{g/ml}$ respectively. Result suggests a close relations in between the ethanol extracts of leaves and stem of *G.Montana* antioxidant activity. Therefore it was concluded that ethanol extract of leaves and stem of *G.Montana* showed potent antioxidant activity. It can be used as antioxidant for the treatment of various diseases.

Key Words : - *Gymnospora Montana*, free radical, DPPH, antioxidant, ascorbic acid.

1. INTRODUCTION :-

Antioxidant are the compounds that neutralize free radicals which otherwise damages the body. Free radicals are chemically active product of metabolism and include reactive oxygen species or reactive nitrogen species. Free radicals or oxidative injury now appears the fundamental mechanism underlying a number of human neurologic and other disorders [1]. Various pathological conditions such as cancer, cataracts, chronic inflammation as well as diabetes mellitus, cardiovascular and neurodegenerative diseases believed to be associated with oxidative stress [2,3,4].

Synthetic antioxidants has been widely used. The incorporation of these synthetic antioxidants in food preparations have been questioned due to potential health risks, toxicity and



carcinogenicity [5,6]. It is increasingly being realized that a majority of the disease of today are due to the shift in the balance of the pro-oxidant and the antioxidant homeostatic phenomenon in the body. Pro-oxidant conditions dominate either due to the increased generation of the free radicals or due to the excessive oxidative stress of the depletion of the dietary antioxidant [7]. Plant material have evolved a wide range of mechanisms to contend with this problem, with variety of antioxidant molecules and enzymes.

In recent years much attention has been devoted to natural antioxidant and their association with health benefits and many plants exhaustively studied in the last few years for their antioxidant and radical scavenging activities [8]. *G.Montanaplant* has been used traditionally and useful in treating ulcer, gastrointestinal disorders, toothache, dysentery [9], antispasmodic [10] and hepatoprotective [11] effects were also reported. A detailed review of literature afforded no information on the in-vitro antioxidant potential of the plant were investigated by using DPPH scavenging method.

Therefore, taking into consideration the vast potentiality of plants as sources for antioxidants a systematic investigation was undertaken to screen the local flora for radical scavenging activity. The leaves and stem of *G.Montanawere* extracted in ethanol and tested free radical scavenging activity against 2,2-diphenyl-1-picrylhydrazyl (DPPH). Ascorbic acid was used as antioxidant reference compound.

2. MATERIAL AND METHODS :-

2.1 Plant material:The leaves and stem of *G.Montanawere* collected from Balaghat region of Beed district of Maharashtra.

2.2 Extraction :The leaves and stem of *G.Montanawere* dried under shade and then powdered. The dried powdered material was subjected to extraction with ethanol. 10 gm of powdered plant material was dissolved in 100 ml of ethanol and kept on a magnetic stirrer for 24 hrs. Thereafter, it was filtered and centrifuged at 5000 rpm for 15 min. The supernatant was collected and the solvent was evaporated out to dryness. The obtained material was stored at 4°C in airtight bottles for further studies.



2.3 Evaluation of antioxidant activity :-

DPPH Method

DPPH scavenging activity was measured by the spectrophotometric method. A stock solution of DPPH (1.5 mg/ml in methanol) was prepared such that 75 μ l of it in 3 ml of methanol. Decrease in the absorbance in the presence of sample extract at different concentration (10-125 μ g/ml) was noted after 15 min. IC₅₀ value was calculated from % inhibition.

Preparation of stock solution of the sample :-

10 mg of extract was dissolved in 10 ml of methanol to get 1000 μ g/ml solution. Then were prepared diluted test solution of different concentration as 10, 20, 30, 40, 50, 60, 70, 80, 90 and 100 μ g/ml from stock solution. 15 mg for DPPH solution was dissolved in 10 ml of methanol. The resulting solution was covered and protected from light.

75 μ l of DPPH solution was taken and the final volume was adjusted to 3 ml with methanol, absorbance was taken immediately at 517 nm for control reading. 75 μ l of DPPH and 100 μ l of the test sample of different concentration were put in a series of volumetric flasks and final volume was adjusted to 3 ml with methanol. Absorbance at zero time was taken in UV-visible spectrophotometer at 517 nm for each concentration. Final decrease in absorbance of DPPH with sample of different concentration was measured after 15 minute at 517 nm.

Percentage inhibitions of DPPH radical by test compound were determined by the following formula.

$$\% \text{ Inhibitions} = \left(\frac{\text{Control absorbance} - \text{Test absorbance}}{\text{Control absorbance}} \right) \times 100$$

IC₅₀ value was calculated by using graphical method.

3. OBSERVATION :-

Antioxidant activity testing



Table.1. DPPH Free Radical Scavenging Activity of Ascorbic Acid.

Absorbance of the sample at 517nm Absorbance of Control = 0.490

Sr. No.	Concentration In (µg/ml)	Absorbance of Ascorbic acid	% Inhibition	IC ₅₀ Value (µg/ml)
1	10	0.295	39.79	25
2	20	0.270	44.89	
3	30	0.246	49.79	
4	40	0.225	54.08	
5	50	0.203	58.57	
6	60	0.184	62.44	
7	70	0.161	67.14	
8	80	0.142	71.02	
9	90	0.120	75.57	
10	100	0.087	82.24	

Table.2. Antioxidant Activity of Ethanol Extract of *G.Montana*(leaves)

Absorbance of the sample at 517nm Absorbance of Control = 0.494

Sr. No.	Concentration In (µg/ml)	Absorbance of Ascorbic acid	% Inhibition	IC ₅₀ Value (µg/ml)
1	10	0.289	41.49	34
2	20	0.273	44.73	
3	30	0.261	47.16	
4	40	0.249	49.59	
5	50	0.232	53.03	
6	60	0.219	55.66	
7	70	0.196	60.32	
8	80	0.181	63.36	
9	90	0.165	66.59	
10	100	0.156	68.42	



Table.3. Antioxidant Activity of Ethanol Extract of *G.Montana*(stem)


Absorbance of the sample at 517nm Absorbance of Control = 0.494

Sr. No.	Concentration In (µg/ml)	Absorbance of Ascorbic acid	% Inhibition	IC ₅₀ Value (µg/ml)
1	10	0.292	40.89	36
2	20	0.275	44.33	
3	30	0.268	45.74	
4	40	0.253	48.78	
5	50	0.237	52.02	
6	60	0.221	55.26	
7	70	0.210	57.48	
8	80	0.196	60.32	
9	90	0.183	62.95	
10	100	0.177	64.17	

4. RESULTS AND DISCUSSION :-

In vitro antioxidant activity of ethanol extracts of leaves and stem of *G.Montana* were tested for their antioxidant activity of DPPH scavenging method. Result shows that it has better antioxidant activity. It is observed that, phenolic compounds are responsible for antioxidant activity [12,13].

The phenolic compounds from *G.Montana* may be responsible for antioxidant activity. Phenolic compounds are effective hydrogen donors, which make them antioxidant [14]. The dose response curve of DPPH for extracts of *G.Montana* leaves and stem was compared with reference standard ascorbic acid as shown in figure.



DPPH scavenging activity was found in ethanolic extracts of *G. Montana* leaves and stem. Compared to ascorbic acid the radical scavenging activities of both test samples has their IC_{50} values ranged from 34 and 36 $\mu\text{g/ml}$ respectively in DPPH assay, (Table-2,3).

This indicates that ethanolic extracts of *G. Montana* leaves and stem has good potential as a source for natural antioxidants. Ascorbic acid was used as a standard. Which shows IC_{50} value of 25 $\mu\text{g/ml}$ in DPPH assay, (Table-1).

5. CONCLUSION :-

The results obtained in present study were indicated that extracts of *G. montana* inhibits free radical scavenging activity. The overall antioxidant activity depends on its triterpenoid and polyphenolic content and other phytochemical constituents were present. It could be a source of natural antioxidant that could have greater importance as therapeutic agent in preventing or slowing oxidative stress related degenerative diseases.

It can be observed that ethanolic extract of leaves and stem of *G. montana* showed very close significant scavenging activity as compared to the reference standard ascorbic acid.



Therefore, it was concluded that ethanolic extract of *G. Montana* showed potent antioxidant activity.

ACKNOWLEDGMENT :-

Authors are thankful to Principal and Head of Department of our college for motivation, support and providing necessary facilities to carry out study.

REFERENCES :-

1. Nadkarni, K.M. Indian Materia Medica, 3rd Edn, Dhootapapeshwar Prakashan, Ltd., Bombay.1954, 1; 353.
2. Pelicano H, Carney D, Huang P; ROS stress in cancer cells and therapeutic implications, *Drug Resist Update* 2004, 7, 97-110.
3. Gonçalves C, Dinis T, Batista M T; Antioxidant properties of proanthocyanidins of *Uncaria tomentosa* bark decoction: a mechanism for anti-inflammatory activity. *Phytochemistry* 2005, 66, 89-98.
4. Lim Y Y, Murtijaya J, Antioxidant properties of *Phyllanthus amarus* extracts as affected by different drying methods. *LWT-Food Sci Technol* 2007, 40,1664-1669.
5. Jeong S, Kim S, Kim D, Jo S, Nam K, Ahn D, Lee S. 2004.Effect of heat treatment on the antioxidant activity of extracts from citrus peels. *J. Agric. Food Chem.*2006, 52, 3389-3393.
6. Wong S.P, Leong L.P, Koh J H W; Antioxidant activities of aqueous extracts of selected plants. *Food Chem* 99:775-783.
7. Priyesh, P., Narayanasamy, V.B., Manjunatha, Setty, M., Annie, S. Antioxidant potential of *Clerodendron viscosum vent.* Roots. *Pharmacologyonline* 2; 2007, 226-235.
8. Arnous A, Makris D.P, Kefalas P; Effect of principal polyphenolic components in relation to antioxidant characteristics of aged red wines. *J. Agric. Food. Chem*, 2001, 49, 5736-5742.
9. Kirtikar, Basu, B.D. Indian Medicinal Plants International book distributors, Book seller and publisher, Rajpur road, Dehradun, India. 1; 1993, 577.
10. De, S., Dave, K.K., Bhavsar, G.C. Phytochemical study of *Gymnosporia Montana*. *Indian J Pharma Sci.* 55; 1993, 110-112.



11. Auddy, B., Ferreira, M., Blasina, F., Lafon, L., Arredondo, F., Dajas, F., Tripathi, P. C., Seal, T., Mukherjee, B. Screening of antioxidant activity of three Indian medicinal medicinal plants, traditionally used for the management of neurodegenerative diseases. *J. Ethanopharmacol.* 84; 2003, 132.
12. Vijay C, Ramanathan M, Subburaj T and Sures B: Correlation of Phenolic Content and in Vitro Antioxidant Activity of certain herbal extracts, *Indian Drugs* 2002; 3: 453-455.
13. Datir SB, Nirmal SA, Ganjare AB, Bhawar SB and Patil MJ: Antioxidant Activity of the Aerial Parts of the *Achyranthes aspera* Var. *Porphyristachya* (Wall. Ex Moq.) Hook. F. *Research Journal of Pharmacognosy and Phytochemistry* 2009; 1(3): 220-223.
14. Rice-Evans CA, Miller NJ, Bolwell PG, Bramley PM and Pridham JB, Therelative antioxidant activity of plantderived polyphenolic flavonoids. *FreeRadic. Res*, 22: 375 – 383, (1995).