



Available online at <http://www.jmcd.com>

November-December, 2015, Vol. 4, pp 01-13

ISSN: 2347-9027

---

## Research Article

---

### **Assessment of effects of extract from roots and leaves of *Citrullus lanatus*, (THUNB)**

**Popat Ritesh R\***, Mhaske Shivshankar D, Wadekar Ashish B; Narkhede Mahesh B, Shrikhande Vinayak N

IBSS College of Pharmacy, Malkapur

\* Corresponding Author: Email: [riteshpopat87@gmail.com](mailto:riteshpopat87@gmail.com)

#### **Abstract**

*Citrullus lanatus* Thunb (Family: Cucurbitaceae) is a medicinal plant widely used traditionally in the treatment of various disorders in world. The phytochemical evaluation of plant showed presence of carbohydrate, alkaloids, steroids, saponins, glycoside, flavonoids, tannins and phenolic compounds. The plant also showed presence of vitamins, amino acids, proteins, minerals and fat. The plant has been extensively studied by various scientists and researchers for its pharmacological activities and therapeutic approaches such as antibacterial, antifungal, antimicrobial, antiulcer, antioxidant, anti-inflammatory, gastroprotective, analgesic, laxative, anti-giardial, hepatoprotective, against prostetic hyperplasia and atherosclerosis. The present review is an effort to provide detailed information of its uses, chemical composition, pharmacological activities of extract and its isolated compounds and safety profile of *Citrullus lanatus* for further research.



# Journal of Medicinal Chemistry and Drug Discovery

## Introduction

Cucurbitaceae is the largest family containing 120 genera and approximately 825 species typically distributed in the tropical countries poorly represented in the temperate regions. The vegetable crops of family Cucurbitaceae are important horticultural crop, mostly grown for its sweet and juicy fruit in warm climates all over the world. Cucurbitaceae are important source of food like pumpkin (*Cucurbita pepo*), melon (*Cucumis melo*), cucumber (*Cucumis sativa*), water melon (*Citrullus lanatus*), *Lag enaria siceraria* (bottle gourd), and *Luffa cylindrica* (sponge gourd). The *Citrullus lanatus* L is one of the most popular species with high water content as high as 92% of the total weight<sup>1</sup>. The plant is traditionally used for centuries in the treatment of various health ailments. It is an important medicinal plant in the Ayurveda and Indian traditional system of medicine<sup>2</sup>. The plant is rich in flavonoids, alkaloids, saponins, glycoside, tannins and phenols. Its nutritive values are also useful to the human health. The plant has been extensively studied by various scientists and researchers for its pharmacological activities and therapeutic approaches such as antibacterial, antifungal, antimicrobial, antiulcer, antioxidant, anti-inflammatory, gastroprotective, analgesic, laxative, anti-giardial, hepatoprotective and against prosthetic hyperplasia and atherosclerosis. Fruit is used in cooling, strengthening, aphrodisiac and astringent to the bowels, indigestible, expectorant, diuretic, and stomachic, blood purifier. It also allays thirst, cures biliousness, good for sore eyes, scabies and itches and as brain tonic to the brain<sup>1</sup>. The present review focuses on phytochemical, pharmacognostic, pharmacological and toxicity updates of this plant.

## Botanical Description:

### Taxonomy:

Botanical name: *Citrullus lanatus* (Thunb)

Class: Equisetopsida

Kingdom: Plantae

Genus: *Citrullus*

Family: Cucurbitaceae

## Journal of Medicinal Chemistry and Drug Discovery

Order: Cucurbitales

**Vernacular names:**

Common name: Watermelon, Wild Watermelon

Local name: Tarbooz

English: Watermelon

Marathi: Tarbooz, Kalingad

Bengali: Tormuz


Malayalam: Thannimathan

Kanada: Kallagadi

Assamese: Tarmuj

Telugu: Pendalam

Tamil: Kizhangu





# Journal of Medicinal Chemistry and Drug Discovery

## Morphological characters:

*Citrullus lanatus* is a prostrate or climbing annual with several herbaceous, rather firm and stout stems up to 3 m long; the young parts are densely woolly with yellowish to brownish hairs while the older parts become hairless. The leaves are simple, alternate on long petioles, cordate with seven shallow lobes and variously serrated margins, very hairy on the abaxial surface, acute, deep green, and about 7 – 15 cm in diameter. Tendrils are simple and spiral. Male and female flowers grow on the same plant. Male flowers are found in clusters and appear before the female flowers. Both have yellow petals, five in number, and sepals, also five in number and greenish in color. Occasional hermaphrodite flowers are produced. The fruits are globular with shallow grooves, about 14 – 20 cm long. The skin is greenish yellow. The flesh is almost white/light yellow, sweet, delicately flavoured, juicy, a pepo. The seeds are small, light brown and smooth, between 0.4 and 1.1 cm long and 0.2 – 0.3 cm wide.

## Traditional claims:

*Citrullus lanatus* (Thunb) is used as anthelmintic, anticancer, antibacterial, demulcent and diuretic being used in the treatment of dropsy and renal stones. The rind of fruit is prescribed in case of alcoholic poisoning and diabetes. The root is purgative and in large doses emetic. The seed is demulcent, diuretic, pectoral and tonic. It is sometimes used in treatment of bed wetting. Seeds are also vermifuge and have hypotensive action. Fatty oils in the seeds as well as in aqueous and alcoholic extracts paralyze tapeworms and roundworms. In Northern Sudan, it is often used for burns, swellings, rheumatism, gout and as laxative. Fruits of this plant are eaten as febrifuge when fully ripe or when almost putrid. It is also used in case of energy source, cleanses and purifies kidney and bladder, prevent erectile dysfunction and used to treat hepatomegaly and jaundice.

## Chemical Constituents:

*Citrullus lanatus* seed contains phytochemical constituents like alkaloids, flavanoids, tannins, amino acids, carbohydrates, cardioglycosides, terpenoids, steroids, carotenoids, oils and fats. The amino acid citrulline was first extracted from watermelon and analyzed. The nutritional quality of watermelon shows that it is very rich in vitamins A 3%, different vitamins from vitamin B complex like Thiamine (Vit. B1), Riboflavin (Vit. B2), Niacin (Vit. B3), Pantothenic acid (B5), vitamin B6 and Folate (Vit. B9) which ranges between 1-3%, Vitamin C 14%. The

## Journal of Medicinal Chemistry and Drug Discovery

mineral composition is Calcium 1%, Iron 2%, Magnesium 3%, Phosphorus 2%, Potassium 2% and Zinc 1%. Along with it, it also contains highly unsaturated fatty acids and oils. It is also rich in essential amino acids like arginine, glutamine and aspartic acid 8.


### Pharmacology:

#### Antibacterial activity and antifungal activity:

Crude extract of watermelon seeds using hot water, cold water, methanol and ethanol showed the antimicrobial activity using the standard disc diffusion assay method against *Staphylococcus aureus*, *Pseudomonas aeruginosa*, *Escherichia coli*, *Klebsiella pneumoniae* and *Bacillus cereus*. All these seed extracts showed evidence of antibacterial properties 9. In another study, crude extract of *Citrullus colocynthis*, *Citrullus lanatus* and *Citrullus vulgaris* were very effective against bacteria and some fungal strains. The ethanolic extract was found more effective than respective aqueous and chloroform extract. The zone of inhibition was found maximum in *E. coli* and minimum in *Staphylococcus aureus*. Fungal zone of inhibition was found more in *Candida albicans* and less in *Trichosporon Begelii* 10.


#### Figure-2. Amino acids in *Citrullus lanatus*

(a-Citrulline, b-Glutamine and c-Aspartic acid)




(a) Citrulline

(b) Glutamine



## Journal of Medicinal Chemistry and Drug Discovery

### (C) Aspartic acid



### Antimicrobial activity:

The antimicrobial effects of chloroform of *Citrulluslanatus* extracts against the studied bacteria suggestthat, different parts of plant possess remarkabletherapeutic action that can support the traditionalusage of this plant in the treatment of bacterial andfungal diseases such as gastrointestinal infection,diarrhea, respiratory and skin diseases. Theseantimicrobial activities are likely due to the presenceof secondary metabolites like tannins and flavonoids,alkaloids, saponins, terpenes and steroids in plant.The high potency of *Citrullus lanatus* against thesemicrobes could provide an example of prospectingfor new compounds 6. In another study, antimicrobialefficacy of *Citrullus lanatus* and other citrullus plantswere evaluated against gram negative, positive andsome fungal strains including *E-coli*, *Klebsiellapneumonia*, *Pseudomonas aeruginosa*, *Bacillussubtilis* and *Bacillus pumilus* some fungal *Candidaalbicans*, *Aspergillus niger*, *Penicillium chrysogenum*using agar well diffusion method at concentration 100mg/mL. In this, almost all species of plants werefound to have activity on at least two microbial strains7. Determination of antimicrobial activity of methanolextract of *Citrullus lanatus* seed was carried outagainst 10 bacterial species (*Staphylococcus aureus*,*Klebsiella pneumoniae*, *Bacillus subtilis*, *Escherichiacoli*, *Salmonella typhi*, *Enterococcus faecalis*, *Vibriocholerae*, *Shigella dysenteriae*, *Proteus mirabilis*,*Pseudomonas aeruginosa*) and 5 fungal species(*Aspergillus flavus*, *Aspergillus niger*, *Penicilliumnotatum*, *Trichophyton mentagrophytes*, *Candidaalbicans*). Inhibition zones formed by the extract werecompared with the standards: Streptomycin andAmphotericin B. The methanolic seed extract at aconcentration of 1000 µg/ml was effective againstbacteria: *Vibrio cholerae* (7 mm), *Proteus mirabilis* (7mm), *Shigella dysenteriae* (7 mm), *Staphylococcus aureus* (6 mm), *Escherichia coli* (6 mm),*Enterococcus faecalis* (6 mm), *B. subtilis* (6 mm),*Salmonella typhi* (5 mm) and fungi: *Aspergillus niger*(13 mm), *Aspergillus flavus* (12 mm),



## Journal of Medicinal Chemistry and Drug Discovery

*Penicillium notatum* (11 mm), *Candida albicans* (9 mm). The results of *Citrullus lanatus* seed extract demonstrated antimicrobial activity against the organisms tested 11.

### **Antiulcerative activity:**

Methanolic extract of *Citrullus lanatus* (MECL) seeds showed maximum antioxidant potential and was evaluated for its anti-ulcerogenic activity by Pyloric Ligated (PL) and Water Immersion Stress (WIS) induced ulcer models in rat. Gastric volume and free and total acidity were measured in PL model whereas; ulcerative index was measured in both the models at 100, 200 and 300 mg/kg doses of MECL. Rats treated with MECL (300 mg/kg) showed significant decrease in the gastric volume, free acidity and total acidity in case of PL model and showed significant percentage inhibition of ulcer as indicated by decrease in ulcerative index in both the models. The extract of *Citrullus lanatus* seeds possesses good antioxidant and anti-ulcer activity 12. In another study, the presence of secondary metabolites such as flavonoids, tannins, saponins, terpenoids claimed to present in the most members of the Cucurbitaceae family. Proximate and metal content analysis of these seeds provides information that the consumption of the seeds of *Citrullus lanatus* is safe. This study also provides preliminary data for the first time that these seeds of *Citrullus lanatus* possess significant antiulcer activity in animal model. The anti-ulcer activity is probably due to the presence of bioactive compounds like flavanoids, saponin and tannins. Further studies are required to confirm the exact mechanism underlining the ulcer healing and protecting property of the extract and to identify the chemical constituents responsible for it 13. The antiulcer activity of crude methanolic extract of *Citrullus lanatus* seeds in two different ulcer models in albino Wistar rats was also studied. The extract at 300 mg/kg body weight, once daily orally for 7 days has a significant effect in pyloric ligation (PL, 4 h ligation) and in water immersion (WS, 25 °C for 3 h), stress induced ulcer model, as it showed protection index of 57.33% and 63.38% respectively which is comparable to the standard drugs (Ranitidine 50 mg/kg) and Omeperazole (20 mg/kg body weight), that have shown protection index of 64.47% and 70.59% in PL and WS model respectively 14.

### **Antioxidant activity:**



## Journal of Medicinal Chemistry and Drug Discovery

The In-vitro anti-oxidant activity of the n-Hexane, Chloroform and Ethanol extract of *Citrullus lanatus* seeds were studied using 1,1-diphenyl-2-picrylhydrazyl (DPPH) radical scavenging activity, Ferric reducing power activity, Hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>) scavenging activity and Nitric Oxide (NO) scavenging activity. The total Phenolic contents and Flavonoid contents were estimated taking Gallic Acid and Quercetin calibration curve respectively. In this study, it was found that all the extracts possess In-vitro antioxidant activities. But the order of possessing activities were n-hexane > ethanol > chloroform extracts of *Citrullus lanatus* seeds 15.

### **Anti-inflammatory activity:**

*Citrullus lanatus* seed oil is extracted with n-hexane and tested for in-vivo and in-vitro anti-inflammatory activity. The oil was screened for in-vivo anti-inflammatory activity by Carrageenan-induced paw edema in rat model and In-vitro anti-inflammatory activity by human red blood cell membrane stabilization method. The potency of the oil was compared with standard Diclofenac (10 mg/kg). The oil showed significant reduction of edema in Carrageenan induced Rat Paw Edema Model maximum at 3 hr (percentage reduction in paw volume 44.44%, 55.56% and 63.11% for CLSO (50 mg/kg), CLSO (100 mg/kg) and diclofenac (10 mg/kg) respectively and CLSO at concentration of 100, 250 and 500 mcg/ml showed 42.35%, 68.48% and 78.50% protection of human red blood cell membrane in hypotonic solution respectively 16. In another study, mouse model with ear edema induced by xylene and the rat model with paw edema or granuloma by carrageenan or cotton pellet were used for anti-inflammatory effect of the *Citrullus lanatus* aqueous extract. Effects of the extract on analgesia were tested respectively by measuring the latency of mice licking hind foot from hot plates and by counting the times of body twisting in response to acetic acid. The extract significantly inhibited the ear edema, granuloma hyperplasia and paw edema. It significantly lifted pain threshold on mouse hot-plate responses and reduced their writhing time 17.

### **Gastroprotective Activity:**

Gastroprotective potential of *Citrullus lanatus* fruit pulp aqueous extract (CLE) on pyloric ligation and indomethacin induced ulcer model in wistar albino rats was evaluated. In indomethacin induced ulcer model, CLE was administered in the doses of 250 mg/kg, 500 mg/kg & 1000 mg/kg body weight orally for five days for





## Journal of Medicinal Chemistry and Drug Discovery

three times in a day after the induction of ulcer. The extent of healing was determined via reduction in ulcer index and alteration in various biochemical parameters in gastric mucosa. In pyloric ligation model, CLE resulted in significant increase in pH, enzymic antioxidants i.e. Superoxide dismutase & Catalase, with a significant decrease in volume of gastric juice, free and total acidity, protein & carbohydrate concentration and Lipid peroxidase levels. The presence of the flavonoids and polyphenols may be responsible for the gastroprotective effect of CLE 18.

### Activity against prosthetic hyperplasia:

Administration of MECLS for one month reduced the prostate size significantly ( $P < 0.05$ ), both at high and low dose, but could not restore the initial size of shrunken testes and severe oligospermia caused by the hormones. The histological studies clearly establish MECLS as a potential candidate in management of androgen dependent conditions like benign prostate hyperplasia 8.

### Laxative Activity:

The laxative activity was determined based on the weight of the fecal matter. The effects of the aqueous fruit pulp extract of *Citrullus lanatus* and reference standard on the gastro intestinal motility rate were also evaluated. Phytochemicals screening of the extract revealed the presence of flavonoids, tannins, polyphenols, sterols and polyterpenes. The aqueous fruit pulp extract of *Citrullus lanatus* administered orally at three different doses produced significant laxative activity and reduced loperamide induced constipation in dose dependent manner. The effect of the extract at 500 and 1000 mg/kg (p.o.) was similar to that of reference drug sodium picosulfate (5 mg/kg, p.o.). The same doses of the extract (500 and 1000 mg/kg, p.o.) produced a significant increase ( $p < 0.01$ ) of intestinal transit in comparison with castor oil (2 ml) ( $p < 0.01$ ). The results showed that the aqueous fruit pulp extract of *Citrullus lanatus* has a significant laxative activity 19.

### Antigiardial Activity:

Antigiardial activities of *Citrullus lanatus* var. *citroides* (wild watermelon) fruits, petroleum ether, ethyl acetate, butanol crude extracts as well as Cucurbitacin E and Cucurbitacin L 2-O- $\beta$ -glucoside pure isolated compounds from *Citrullus lanatus* var. *citroides* was investigated. In this study, Cucurbitacin E and Cucurbitacin L 2-O- $\beta$ -glucoside were revealed to have strong potent anti-giardial activity against *Giardia lamblia* in vitro with  $IC_{50} = 2$



## Journal of Medicinal Chemistry and Drug Discovery

and 5 ng/ml after 5 days respectively. The ethyleacetate extract was the best among all examined extracts followed by petroleum ether and butanol with IC<sub>50</sub> 0.1, 0.2 and 0.5 µg/ml respectively. The results suggested that all the crude extracts and isolated compounds were reactive against *G. lamblia*, hence *Citrullus lanatus* var. *citroides* may be recommended as a new source for the treatment of giardiasis 20.

### **Anti-hepatotoxic activity:**

In the study, the ability of watermelon juice to protect against CCl<sub>4</sub>-induced hepatotoxicity and oxidative stress was investigated. It has been hypothesised that one of the principal causes of CCl<sub>4</sub>-induced liver injury is formation of lipid peroxides by free radical derivatives of CCl<sub>4</sub>. The trichloromethyl radical produced by carbon tetrachloride leads to liver damage by alkylating cellular proteins and other macromolecules with a simultaneous attack on polyunsaturated fatty acids to produce lipid peroxides (Lipid peroxidation, a ROS-mediated mechanism, has been implicated in the pathogenesis of various liver injuries and subsequent liver fibrogenesis in experimental animals. Hepatocellular necrosis leads to elevations of serum AST and ALT activities and an increased incidence and severity of histopathological hepatic lesions in rats 21.

### **Anti-atherosclerotic activity:**

Administration of *C. lanatus* 'sentinel' extract attenuated atherosclerosis in both arch and thoracic regions of aortas in mice. Aortic arch and thoracic aortic regions represent different developmental origins of aortic smooth muscle cells and reduction of atherosclerosis in these two distinct aortic regions provides strong evidence of a role for *C. lanatus* 'sentinel' extract. There is evidence that the main component of *C. lanatus*, citrulline, has a beneficial effect on atherosclerosis and aortic vascular remodeling. Citrulline administration improved endothelium-dependent vasorelaxation and decreased aortic medial thickening. It is possible that citrulline in *C. lanatus* 'sentinel' extract may contribute to reducing atherosclerosis 22.

### **Anti-secretory activity:**

The effects of the juice of *Citrullus lanatus* (watermelon) was evaluated on gastric acid secretion and pH in Indomethacin-induced ulceration in male albino rats. Rats pre-treated with *Citrullus lanatus* juice exhibited



## Journal of Medicinal Chemistry and Drug Discovery

significant dose-dependent reduction of gastric lesions formation ( $P < 0.05$ ). Also, ulcerogenesis in the pretreated groups was significantly lower than that observed with the control ( $P < 0.05$ ) 23.

### **Analgesic activity:**

The analgesic activity of aqueous extract of *Citrullus lanatus* peels (AECL) using Eddy's hot plate method was evaluated. The AECL produced a significant analgesic activity in a dose dependent manner. All the doses of AECL (250, 500 and 1000 mg/kg) had shown a good analgesic activity. The reaction time obtained for these three doses after 90 minutes of drug administration was found out to be 5.15 mins, 8.92 mins and 10.82 mins respectively which was comparable to Diclofenac sodium (5 mg/kg) that showed the reaction time of 12.36 mins 24.

### **Safety and toxicity profile:**

Acute toxicity of *Citrullus lanatus* extracts were studied in rodents. The n-hexane extract of *Citrullus lanatus* seed oil (CLSO) using acute toxic class method was evaluated as described in OECD guidelines no. 423. The *Citrullus lanatus* seed oil was found safe up to a dose of 2,000 mg/kg body weight [16]. The toxicity study of aqueous extract of roots and leaves was also evaluated on mice. No mice died during the observation period so maximum tolerance experiment was carried out according to GLP 2003 by state food and drug administration. Twenty mice were given orally single maximum dose of 43.5 g/kg at two time points (7 am and 5 pm) but no mice died in period of 7 days [17]. The ethanolic extract of *Citrullus lanatus* was also found to show no mortality up to the maximum dose level of 2000 mg/kg body weight of the extract after administered orally 25.

### **Conclusion**

*Citrullus lanatus* (Thunb.) has been indicated in the treatment of various health ailments by traditional and Ayurvedic system of medicine. The research carried out till now have confirmed pharmacological potential of *Citrullus lanatus* and is found to be safe. Further research is necessary to reveal its detailed molecular mechanism behind these pharmacological activities.

### **Conflict of Interests**

The authors declare that there is no conflict of interests regarding the publication of this paper.



# Journal of Medicinal Chemistry and Drug Discovery

## References

1. Mabberley DI, 1987. The Plant Book. Cambridge:Cambridge Univ. Press, New York.
2. Gill NS, Kaur S, Arora R, Screening of antioxidant and anti-ulcer potential of *Citrullus lanatus* methanolic seed extract, Research J of Phytochem 2011; 5 (2): 98-106.
3. Mercy GA, Bosa EO, The morphological characterization of the melon species in the family cucurbitaceae juss., and their utilization in Nigeria, International J of Modern Botany 2013;3(2): 15-19.
4. Jain S, Jerald EE, Edwin S, Isolation, fractionation and evaluation of the anti-inflammatory properties of *Citrullus lanatus* Thunb, Asian J of Biomed & Pharma Sciences, 2013; 3(20):66-72.
5. Varghese S, Narmadha R, Gomathi D, Phytochemical screening and HPTLC fingerprinting analysis of *Citrullus lanatus* (Thunb.) seed, J of acute Disease 2013:122-126.
6. Ahmed Hassan LE, Sirat HM, Yagi SM, In vitro antimicrobial activities of chloroformic hexane and ethanolic extracts of *Citrullus lanatus* var. *Citroides* (wild melon), J of Med Plants Res 2011; 5(8): 1338-1344.
7. Thirunavukkarasu P, Ramanathan T, Screening of antimicrobial effects in watermelon, J of BiolSci 2010:1-4.
8. Olamide AA, Olayemi OO, Demetrius OO, Effects of methanolic extract of *Citrullus lanatus* seed on experimentally induced prostatic hyperplasia, Eur J of Med Pla 2011; 1(4): 171-179.
9. Braide W, Odiong II, Oranusi S, Phytochemical and antibacterial properties of the seed of watermelon (*Citrullus Lanatus*), Prime J of MicroRes 2012; 2(3): 99-104.
10. Thirunavukkarasu P, Ramanathan T, Screening of antimicrobial effects in watermelon, J of BiolSci 2010: 1-4.
11. Sathya J, Shoba FG, Assessment of antimicrobial efficacy of *Citrullus lanatus* methanolic seed extract, J. Chem. Pharm. Res 2014; 6(12):640-643.
12. Gill NS, Sood S, Evaluation of antioxidant and anti-ulcerative potential of *Citrullus lanatus* seed extract in rats, Lat. Am. J. Pharm 2011; 30 (3):429-434.
13. Lucky OO, John UO, Quantitative determination, metal analysis and antiulcer evaluation of methanol seeds extract of *Citrullus Lanatus* thunb (Cucurbitaceae) in Rats, Asian Pac J of Trop Dis 2012; S1261-S1265.



## Journal of Medicinal Chemistry and Drug Discovery

14. Bhardwaj A, Kumar R, Dabasa V, Evaluation of anti-ulcer activity of *Citrullus lanatus* seed extract in wistar albino rats, *Int J Pharm Pharm Sci*; 4(5):135-139.
15. Rahman H, Manjula K, Anoocha T, In-vitro antioxidant activity of *Citrullus lanatus* seed extracts, *Asian J Pharm Clin Res* 2013; 6 (3):152-157.
16. Madhavi P, Maruthi R, Kamala V, Evaluation of anti-inflammatory activity of *Citrullus lanatus* seed oil by *in-vivo* and *in-vitro* models, *Int. Res J Pharm. App Sci* 2012; 2(4):104108.
17. Deng JG, Wang S, Anti-inflammatory and analgesic effects of extract from roots and leaves of *Citrullus lanatus*, *Chine Herb Med*, 2010; 2(3):231-235.
18. Sharma S, Jain S, Dwivedi J, Gastroprotective activity of *Citrullus lanatus* in rats, *Apac J of Trop Biomed* 2012; 1-6.
19. Sharma S, Paliwal S, Dwivedi J, First report on laxative activity of *Citrullus lanatus*, *Pharmacologyonline* 2011; 2: 790-797.
20. Hassan LE, Koko WS, *In vitro* anti-giardial activity of *Citrullus lanatus* var. *citroides* extracts and cucurbitacins isolated compounds, *J of Med PlaRes* 2011; 5(15): 3338-3346.
21. Altas S, Kizil G, Protective effect of Diyarbakır watermelon juice on carbon tetrachloride-induced toxicity in rats, *Food and Chem Toxicol* 2011;49: 2433–2438.
22. Poduri A, Rateri DL, *Citrullus lanatus* 'sentinel' (watermelon) extract reduces atherosclerosis in LDL receptor-deficient mice, *J of Nutri Biochem* 2013; 24: 882–886.
23. Oluwole FS, Balogun ME Antisecretory effects of watermelon (*Citrullus lanatus*) juice in male albino rats, *Ann Revi & Res in Biol* 2013;3(4): 358-366.
24. Kumari A, Rao J, Kumari J, Analgesic activity of aqueous extract of *Citrullus Lanatus* peels, *Advin Pharmacol and Pharma* 2013;1(3): 135-138.
25. Varghese S, Narmadha R, Evaluation of hypoglycemic effect of ethanolic seed extracts of *Citrullus lanatus*, *J of Phytopharmacol* 2013;2(6): 31-40.